



Estimating bioenergy potentials of common African agricultural residues

Thomsen, Sune Tjalfe; Kádár, Zsófia; Schmidt, Jens Ejbye

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Thomsen, S. T., Kádár, Z., & Schmidt, J. E. (2013). *Estimating bioenergy potentials of common African agricultural residues*. Poster session presented at The 35th Symposium on Biotechnology for Fuels and Chemicals, Portland OR, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Estimating bioenergy potentials of common African agricultural residues

Sune Tjalfe Thomsen^a, Zsófia Kádár^a, Jens Ejbye Schmidt^{a,b}

^asunt@kt.dtu.dk Center for Bioprocess Engineering,

^aDept. of Chemical and Biochemical Engineering, Technical University of Denmark, DK-2800 Lyngby, Denmark

Postal Address: DTU RISØ Campus, Building 301, Frederiksborgvej 399, DK-4000 Roskilde, Denmark

^bPresent address: Chemical Engineering Program, Masdar Institute, PO Box 54224, Abu Dhabi, United Arab Emirates



Introduction


Asking a bioenergy researcher about the composition of wheat straw, he would know it by heart. But if enquiring about typical African biomasses – it would be another case. Until now, biomasses common to African countries have not received the same scientific attention as biomasses from Europe, North America or Brazil. For that reason, it is difficult to estimate bioenergy potentials in the African region.

As a part of an on-going research collaboration investigating production of 2g biofuels in Ghana, this study have analysed 13 common African agricultural residues: yam peelings, cassava peelings, cassava stalks, plantain trunks, plantain leaves, cocoa husks, cocoa pods, maize cobs, rice straw, groundnut straw and oil palm EFB. This was done to establish detailed compositional mass balances, enabling estimations of accurate bioenergy potentials for bioethanol and biogas. Furthermore, biomasses high in specific biomass constituents, such as hemicellulose or lignin, have been identified for future biorefinery applications.




Results


Composition of common African agricultural residues



Potential ethanol from African agricultural residues



Theoretical biogas potential from African agricultural residues



Discussion

The composition of the residues shows significant differences. **Yam-, cassava-, and plantain peelings have very high bioenergy potentials** of up to 41 g bioethanol (100 g TS)⁻¹ and 439 L methane (kg TS)⁻¹ due to a high starch content. **Plantain trunks have the highest amount of cellulose**, 46 g (100 g TS)⁻¹ which can hold potentials in relation to biomaterials as well as bioenergy.

Cocoa pods are unusual rich in lignin, thus poor in carbohydrates and will therefore not be optimal for fermentative processes such as in biogas and bioethanol production. However, this residue might be favorable for biorefinery purposes where a phenol rich starting material is desired.

Cocoa husks and groundnut straw are among the residues with least carbohydrates and therefore they hold low ethanol potentials, however these residues are relatively **high in protein** which makes them more attractive as livestock feed.

The bioenergy potentials are strictly theoretically based on the composition of the residues, thus **biomass recalcitrance are not taking into account**. This might have resulted in over-estimations in the presented results. As part of future studies, laboratory pretreatment and fermentation test of the most attractive residues should be done.

The findings in this study will be linked to results from project partners who have assessed availability of the agricultural residues, possible low tech pretreatment technologies, and sustainability of the agricultural systems.



Acknowledgement

This work was supported with by Danida Fellowship Centre of the Danish Ministry of Foreign Affairs, as a part of the project "2GBIONRG", DFC journal no. 10-018RISØ. If interested visit www.2gbionrg.dk.

Annette Eva Jensen is gratefully thanked for technical assistance.

Presented pictures from www.old.ita.org

*



Methodology

Raw material analysis was done according to standard methods (NREL 2012). Ethanol potentials were estimated stoichiometrically, while the theoretical biogas potentials were estimated using Buswell's formula:

